METHODS

Medical records of 16 dogs treated with lomustine for multicentric T-cell lymphoma were retrospectively reviewed. Ten dogs were treated with a CHOP based protocol (three of which had the inclusion of L-asparaginase initially) and received lomustine as a rescue agent once remission was lost. Six dogs were treated with lomustine as first line therapy for T-cell lymphoma. All dogs received single agent lomustine with a median starting dose of 78.5 mg/m² (range 60–90 mg/m²) every three weeks. Dose adjustment and increased dosage intervals were used when myelosuppression was documented. A total of eight treatments were planned in all dogs. Additional prednisolone was administered in three dogs (starting dose 0.5–1.7 mg/kg q24hrs tapered over four weeks). For those completing therapy, drug-free remission periods were evident. Response to treatment, time to relapse, and survival times for all dogs were documented. Signs of relapse included clinical deterioration and recurrence of lymphadenomegaly and cytological confirmation of lymphoma from aspirates obtained from enlarged lymph nodes where possible.

RESULTS

Of the dogs treated with lomustine as a first line agent (n=6) four dogs had died as a result of their disease with a median time to relapse of 57.5 days (range: 44–105). Two dogs were still alive 628 and 1095 days after starting therapy. The median time to relapse for the dogs treated with CHOP prior to receiving lomustine was 112 days (range: 49–322). Eight of these dogs had a response to lomustine as rescue therapy. Three dogs achieved complete remission, five had a partial response or stable disease and two had no response to therapy with progressive lymph node enlargement. Dogs that responded to lomustine as a rescue agent had a median remission period of 50 days (range: 18–649).

CONCLUSION

Lomustine used first line appears to offer a good long term outcome to a proportion of dogs with T-cell lymphoma, further work is required to investigate this finding.

Spatial clustering of neoplasia in a population of dogs from UK first opinion practice

Stephanie Marlow, Anneliese Stell, Kate English, David Brodbelt

Royal Veterinary College, London, UK

INTRODUCTION

There is limited research into the existence of spatial patterns of neoplasia in the UK general canine population. In addition, until now, there have been few sources of veterinary clinical data that reflect the general population of dogs in the UK and that would therefore enable such investigation. This study aimed to analyse spatial patterns of neoplastic disease in general and mast cell tumours (MCTs) specifically, in the UK first opinion practice-attending canine population, in order to generate hypotheses regarding environmental risk factors for neoplasia.

METHODOLOGY

All data derived from the database of the 'VetCompass' programme (Veterinary Companion Animal Surveillance System); which collects de-identified first opinion practice data directly from electronic patient records from over 450 recruited clinics. A case control study was undertaken to evaluate spatial clustering of neoplasia in general and MCTs. Cases were defined as dogs attending one of the VetCompass clinics

included in this study, with a neoplasm confirmed via histopathology, or cytology reported by a specialist centre (2010–12). MCTs were the most frequently diagnosed potentially malignant neoplasm and were thus taken forward for individual spatial analysis. Partial postcodes for all cases and controls were extracted from the database and their centroid coordinates established via an online Google tool using the mean longitude and latitude of all potential postcodes corresponding to the area of the partial postcode. All mapping was completed in QGIS 2.6.1. Presence of clusters with statistically significant high/low relative risk, were identified via the spatial scan statistic using SaTScan software and a Bernoulli probability model. Statistical significance was set at 5%.

RESULTS

There was evidence of statistically significant clustering for the presence of neoplasia overall and also among the MCTs. An increase in relative risk in and around London was apparent for neoplasia overall with a reduced risk in Cambridgeshire and Oxfordshire. Interestingly, one of those areas of high relative risk identified for neoplasia overall centred in Edgware, Greater London, was in contrast an area of low relative risk for MCTs, while a cluster of high relative risk for MCTs was centred in Hertfordshire.

CONCLUSION

Clustering of canine tumours may reflect differential geographical risk of neoplasia. These data, though preliminary, provide a useful base for more detailed analysis of geographical risk factors for neoplasia within the UK general canine population.

Delivered by BSAVA to:

Guest (guest)

IP: 52.6.47.48

On: Wed, 26 Apr 2023 12:47:23