RESULTS

There was no difference ($P=0.15$) in the relative renal medullary expression of 11β HSD2 between dogs euthanased for behavioural reasons ($n=10$), non-cardiac disease ($n=6$) and CHF ($n=9$). Dogs euthanased for CHF had underlying dilated cardiomyopathy or atrioventricular valve dysplasia.

STATEMENT (CONCLUSIONS)

There is no evidence that modification of expression of 11β HSD2 contributes to the development of canine CHF. However, translational and post-translational modifications of 11β HSD2 and also breakdown of cortisol may be altered, and this warrants further investigation.

Echocardiographic reference intervals in UK Deerhounds

**Emily Dutton¹, Peter Cripps²,
Joanna Dukes-McEwan³**

- 1 Cheshire Cardiology, Knutsford, United Kingdom
- 2 PJC Clinical Epidemiology, Neston, United Kingdom
- 3 University of Liverpool, Neston, United Kingdom

OBJECTIVES

To screen a UK population of clinically healthy Deerhounds to generate breed specific echocardiographic reference ranges for the breed and to identify the prevalence of preclinical DCM in this population.

METHODS

A prospective, longitudinal study of asymptomatic Deerhounds ≥ 4 years of age undergoing auscultation, Doppler echocardiography with ECG and cheek swab sampling.

RESULTS

Deerhounds were classified as affected (preclinical DCM) if they scored >6 according to the ESVC taskforce ($n=13$); and normal (N) if they scored ≤ 3 according to the ESVC taskforce ($n=53$). Twelve hounds were classified as equivocal (4, 5, 6). Seven dogs had arrhythmias, of which 6 were ventricular and 1 was in atrial fibrillation. The 53 N dogs consisted of 26 males and 27 females; median left ventricular internal diameter in diastole value for the male dogs was 55 mm and 51 mm for the females. Median left ventricular internal diameter in systole value for the male N dogs was 40 mm and 37 mm for the N females. Variables of systolic function were assessed in both male and female N hounds, with median Simpson's ejection fraction being 52% and 51% respectively and median fractional shortening being 28% for both sexes. Median end-systolic volume index was 47 ml/m^2 in the female N population and 54 ml/m^2 in the male N population.

STATEMENT (CONCLUSIONS)

Deerhounds have a high prevalence of DCM, ranging from 17% (excluding the equivocal group) to 32% (including the equivocal group). They should be actively screened for disease using echocardiography.

A case-control study of end-stage hypertrophic cardiomyopathy in cats

**Sarah Tayler¹, Lois J Wilkie²,
Virginia Luis Fuentes¹**

- 1 Department of Clinical Science and Services, RVC, University of London AL9 7TA, London, United Kingdom
- 2 Cardiff Cat Clinic, Bridge House, Station Road, Llanishen, Cardiff, Cardiff, United Kingdom

OBJECTIVES

Hypertrophic cardiomyopathy (HCM) is considered end-stage (ES-HCM) when systolic dysfunction develops.

Although widely reported in human literature, only case reports of ES-HCM have been reported in cats. We aimed to compare the clinical characteristics of cats with HCM and ES-HCM.

METHODS

Medical records were searched for cats with HCM, defined as LV wall thickness $\geq 6 \text{ mm}$ without an underlying cause. A subgroup of cats with LV fractional shortening $<30\%$ were classified as ES-HCM, and case matched to cats with 'classic' HCM by breed, sex and age.

RESULTS

Prevalence of ES-HCM was estimated at 5.2% of the HCM population. The majority of cats were Domestic Shorthairs, with British Shorthairs appearing over-represented. Cats with ES-HCM were more likely to present without a murmur ($p<0.001$), with an arrhythmia ($p=0.004$) and in CHF ($p=0.006$). Echocardiographic findings showed cats with ES-HCM were likely to have regional LV wall hypokinesia ($p<0.001$), left atrial enlargement ($p<0.001$) and presence of spontaneous echo contrast/thrombus in the left atrium