Oral presentations

tumours were considered biologically aggressive. C-kit mutations were identified in each patient. Two patients received surgery as main treatment (cases 1 and 3). Case 1 had additional adjuvant prednisolone at the time of tumour recurrence. Case 2 received systemic chemotherapy and brachytherapy, due to tumour location. All patients were alive at the end of the study with a median follow up time of 28 months.

STATEMENT (CONCLUSIONS)

MCT is rarely seen in puppies but needs to be included in the list of differentials. Although an aggressive behaviour may be suspected based on tumour characteristics, a more favourable outcome may be seen in these dogs.

Use of neoadjuvant chemotherapy and radiation therapy for the management of two cases of canine extramedullary oral plasmacytomas

Julie Pradel, Sara Verganti, Davide Berlato

Animal Health Trust, Kentford, United Kingdom

OBJECTIVES

Canine extramedullary oral plasmacytomas are locally aggressive but rarely metastasise. When feasible, wide surgical excision is the treatment of choice and can provide a definitive cure. Radiation therapy has been reported for non-surgical cases, while the use of chemotherapy has been poorly described. This abstract describes a new therapeutic approach for unresectable plasmacytomas combining neoadjuvant chemotherapy and radiotherapy.

METHODS

Retrospective case series including 2 dogs with unresectable oral plasmacytoma affecting the caudal maxilla. Treatment response was assessed according to RECIST criteria.

RESULTS

Case 1 presented with stage II oral multicentric plasmacytomas (T2N0M0). Partial response (70% shrinkage) was achieved with melphalan/prednisolone. Follow-up CT showed complete resolution of the soft tissue component, but residual osteolysis at the level of the larger mass. Because of the multicentric nature of the disease, palliative radiotherapy (36Gy in 6 weekly fractions) was used to control the larger lesion, while melphalan/prednisolone was continued for the other smaller lesions. There was no evidence of tumour progression after 23 months.

Case 2 presented with stage III oral plasmacytoma (T3N0M0). Partial response (67% shrinkage) was achieved with melphalan/prednisolone. As before, follow-up CT showed complete resolution of the soft tissue component, but the osteolysis persisted. The dog underwent a definitive course of radiotherapy (48Gy in 16 daily fractions) and chemotherapy was discontinued. There was no evidence of tumour progression after 12 months.

STATEMENT (CONCLUSIONS)

Neoadjuvant chemotherapy should be considered before irradiating oral plasmacytomas because it could reduce radiation toxicity by allowing smaller targets and sparing sensitive structures such as the eyes.