and ceramides; the IL-8 and TSLP release is increased. Treatment with A97614A1 allowed stressed RHE to recover normal morphology, impermeability, TEWL, IL-8 and TSLP amounts, and close to normal amounts of NMFs and ceramides.

STATEMENT (CONCLUSIONS)

The ingredient A97614A1 was shown to have beneficial effects on the mechanical and immunological skin barrier. These results support the potential interest of incorporating it into topical products.

Two case reports of novel cutaneous mycoses in cats in the UK using molecular identification for rapid and accurate diagnosis

**Nikoleta Makri¹, Gavin Paterson²,
Fiona Gregge³, Catriona Urquhart⁴,
Holly McCluskey⁵, Tim Nuttall¹**

- 1 Hospital for Small Animals, Royal (Dick) School of Veterinary Studies, Edinburgh, United Kingdom
- 2 Easter Bush Pathology, Royal (Dick) School of Veterinary Studies, Edinburgh, United Kingdom
- 3 Aberdeen PDSA Pet Hospital, Aberdeen, United Kingdom
- 4 Abervet, Aberdeen, United Kingdom
- 5 Riverside Veterinary Practice, Livingston, United Kingdom

OBJECTIVES

Phaeohyphomycosis and sporotrichosis are saprophytic mycoses affecting humans and animals. Traumatic inoculation results in cutaneous and subcutaneous lesions that can be locally invasive. Traditional culture based methods can be slow and may not sufficiently classify the fungi. Molecular testing however, can achieve a fast definitive identification.

METHODS

Two cats (cat 1–15-year-old male neutered domestic short hair; cat 2–11-year-old female neutered domestic long hair) developed focal suppurative nodular lesions on the dorsal nose. Both cats had been previously diagnosed with diabetes mellitus and were on insulin therapy. There was no response to broad spectrum antibiotics selected following antimicrobial susceptibility tests. Fungal elements were seen on cytology, although initial culture was negative in cat 2. Subsequent isolates were obtained after overnight incubation of fresh discharge on Sabouraud dextrose agar with chloramphenicol. These were submitted for PCR and sequencing of the ribosomal internal transcribed spacers (ITS) with the primers ITS-F and ITS-R.

RESULTS

PCR and sequencing confirmed phaeohyphomycosis with *Exophiala dermatitidis* (cat 1) and sporotrichosis with *Sporothrix pallida* complex (cat 2). Further sequencing of calmodulin and beta-tubulin partial gene sequences confirmed *Sporothrix humicola*.

STATEMENT (CONCLUSIONS)

These are the first reports of *Exophiala dermatitidis* and *Sporothrix humicola* in cats in the UK. The use of molecular methods provided a rapid and definite identification of the fungi. Clinicians and pathologists need to be aware of the emerging use of molecular techniques to achieve rapid and accurate identification of rare fungal microorganisms. This will help reduce unnecessary antibiotic use.

Antimicrobial susceptibility of bacterial isolates from canine wounds in a referral population

**Catrina Pennington¹, Tim Nuttall²,
Jon Hall³**

- 1 University of Edinburgh, Edinburgh, United Kingdom
- 2 University of Edinburgh, Edinburgh, United Kingdom
- 3 University of Edinburgh, Edinburgh, United Kingdom

OBJECTIVES

To report the prevalence and antimicrobial susceptibility of bacteria isolated from canine wounds in a referral population.

METHODS

Data was collected from clinical records of dogs with wounds cultured between 01/01/2008 and 01/01/2018. Cases with incomplete clinical information or results were excluded. 1 isolate was included where the same bacteria were isolated from multiple samples in a dog.

RESULTS

392 cases (499 samples) were included. 83.5% of cultures were positive; 54.7% isolated a single organism and 28.8% two or more.