Poster presentations

Wild boar injuries in hunting dogs: A retrospective study (2012–2016)

Panagiotis Kokkinos¹, Demetris Tselekis², Georgios Kazakos³, Lysimachos Papazoglou³, Ioannis Tsokatarides⁴

- 1 Animal Health Trust, Lanwades Park, Kentford, Newmarket, Suffolk CB8 7UU, United Kingdom
- 2 Private Practice, Sparti, Greece
- 3 Companion Animal Clinic, School of Veterinary Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece
- 4 Private Practice, Drama, Greece

OBJECTIVES

Description of signalment, wound characteristics, treatment and outcome of wild boar injuries in hunting dogs.

METHODS

In this retrospective study, medical records of hunting dogs from a first opinion and a referral setting were

reviewed for wild boar injuries. Data for 117 dogs were collected and included signalment, physical examination findings, wound characteristics, season of injury, boar sex, type of treatment and outcome among others.

RESULTS

The majority of both the injured dogs (82%) and the boars (85%) were males. Over three quarters of dogs (77%) had a prior history of wild boar injury. The incidents occurred more frequently in December (30%). The most common site of injury was the abdomen (43.6%), followed by the hindlimbs (31.6%) and the thorax (21.4%), whereas 44% suffered multiple injuries. Wound exploration was indicated in 64% of abdominal and 72% of thoracic injuries. Pneumothorax and shock were seen in 20% and 22.4% of dogs, respectively. Surgical treatment was undertaken in 95.6% of the cases. The mean hospitalization time was 4.65 days. The outcome in 95.6% of the cases was good; moreover, 99% of the dogs returned to hunting activity in 26.3 days on average.

STATEMENT (CONCLUSIONS)

The confliction of hunting dogs with wild boars resulted in severe and extended injuries in dogs, usually requiring surgical intervention. Nevertheless, most of the dogs had a good outcome and returned to hunting activity after recovery period. To the authors' knowledge this is the first descriptive study about wild boar injuries in hunting dogs.

Results of artificial urethral sphincter placement in female dogs with refractory urinary incontinence

Joep Timmermans¹, Bart Van Goethem¹, Bouvien Brocks², Hilde De Rooster¹

- 1 Ghent University, Merelbeke, Belgium
- 2 Anicura Dierenkliniek Eersel, Eersel, Netherlands

OBJECTIVES

To determine if placement of an artificial urethral sphincter provides continence in female dogs with refractory urinary incontinence.

METHODS

A retrospective analysis of multi-institutional patient databases resulted in 15 female dogs with refractory urinary incontinence without concurrent underlying disease, treated between September 2010 and March 2018. A hydraulic artificial urethral sphincter with subcutaneous

access port was placed according to the manufacturer's instructions. Treatment success was evaluated using a continence scoring system (scale 1 to 10, complete incontinence to complete continence, respectively) through telephone enquiry.

RESULTS

Median continence score preoperatively was 2/10 (range, 1 to 5). Median follow-up was 6 months (range, 3 months to 5,8 years). Postoperative percutaneous cuff inflation was performed in 7/15 dogs (47%). Complete continence was achieved in 8/15 dogs (53%), improvement in 4/15 dogs (27%), no effect in 2/15 dogs (13%), and worsening in 1/15 dogs (7%). In the patients that improved but were not cured postoperatively, additional medical therapy resulted in a postoperative continence score of 3/10 in 2/5 dogs. Major complications necessitating implant removal surgery occurred in 3/15 dogs (20%), minor complications occurred in 10/15 dogs (67%). Continence scores were awarded to 10/15 dogs (67%). Median continence score postoperatively at follow-up was 9.5/10 (range, 9 to 10).

STATEMENT (CONCLUSIONS)

The artificial urethral sphincter provides promising results in the treatment of bitches with refractory incontinence. In dogs that improve but are not cured after artificial urethral sphincter placement, supplementary medical therapy can further improve the continence rate.