Fracture diagnosis and assessment

Stephen Clarke

Whilst thorough orthopaedic assessment will identify the bone(s) that have been injured, diagnostic imaging is required to classify fracture, facilitate accurate fracture planning and allow for an estimation of costs and prognosis. Sedation or general anaesthesia will be required to facilitate diagnostic imaging and the onus is on the attending clinician to be sure that the traumatized patient has been adequately stabilized beforehand.

Radiography is the most commonly used way of imaging fractures of the appendicular skeleton. A minimum of two high-quality orthogonal radiographs should be obtained, which should include the joints immediately proximal and distal to the fractured bone. For some other regions, for example the carpus and tarsus, oblique and/or stressed projections can provide additional information. Obtaining radiographic projections of the intact, contralateral bone provides a useful resource for fracture planning. Particular attention should be given to the fracture location and configuration as well as the overall appearance of the bones, and these should be considered along with the history and clinical examination findings, in order to be sure that there is no reason, other than trauma, for the development of the fracture.

For certain fractures, advanced imaging will allow superior assessment. Optimizing the information gained may or may not change the eventual management, however it can improve confidence with regards to decision making, aid surgical planning and prognostication. Of the advanced imaging techniques available computed tomography (CT) is the most useful to the fracture surgeon. In the appendicular skeleton CT would be the imaging technique of choice for fractures of the tarsus and carpus; it can also be helpful for assessing pelvic or articular fractures. For fractures of the axial skeleton, CT is invaluable and would be the preferred imaging technique where possible. CT has the added advantage in that it can also allow for rapid prototype modelling for surgical planning and rehearsal, as well as creation of

patient-specific three-dimensional printed drill guides and osteotomy/reduction guides.

KEY LEARNING OBJECTIVES

- Explain the importance of obtaining preoperative radiographs
- Classify fractures based on their location and configuration
- Recognize the situations where advanced imaging may be beneficial

MULTIPLE CHOICE QUESTIONS

1. What does preoperative radiographic fracture assessment allow the surgeon to do?
 - (A) Create a fracture management plan
 - (B) Estimate prognosis for fracture healing and restoration of mobility
 - (C) Provide an estimate of costs
 - (D) All of the above
2. Not all fractures are caused by significant trauma. Which of the following would be the LEAST suggestive that a pathological fracture was present?
 - (A) A comminuted fracture configuration in the absence of significant traumatic injury
 - (B) Mandibular fracture sustained during chewing a toy in a 15-year-old Yorkshire Terrier with renal disease
 - (C) Tibial tuberosity avulsion in a Staffordshire Bull Terrier puppy
 - (D) Folding fracture configuration in an 8-week-old puppy being fed an all-meat diet
3. When assessing a fractured bone, CT can provide superior information than conventional radiography. For which one of these fractures would CT evaluation be of most help?
 - (A) Y-T fracture of the distal humerus
 - (B) Acetabular fracture
 - (C) Comminuted mid-diaphyseal tibial fracture
 - (D) Capital physeal fracture

External skeletal fixation

Eithne Comerford

DEFINITION, PRINCIPLES AND CLASSIFICATION

External skeletal fixators are devices consisting of:

- Percutaneous transcortical pins
- Clamps
- Connecting bars.

External skeletal fixators are classified according to the number and location of connecting bars used: types I–III and hybrid.

IMPLANTS

- Pins. These are placed percutaneously and transosseously and are always bicortical. Pins may be smooth or threaded. Threaded pins have improved pull-out resistance and greater longevity than smooth pins
- Clamps. A number of different makes are available (e.g. Securos, Imex, Veterinary Instrumentation Kirschner Ehmer style). Clamps join pins to the connecting bar(s). Acrylic and epoxy resin can be

Delivered by BSAVA

Guest (guest)

IP: 18.208.226.222

On: Wed, 26 Apr 2023 12:50:32