Triage of polytrauma

Lindsay Kellett-Gregory

Trauma is a common presenting complaint, with a third of trauma cases reported to have serious injuries. All trauma patients therefore require rapid assessment to allow early identification of life-threatening conditions and subsequent stabilization. A traditional major body systems assessment, in which the cardiovascular, respiratory and neurological systems are initially evaluated works well, followed by a secondary screen for any abnormalities elsewhere in the body. Alternatively, the use of a triage scoring system may be useful to identify patients needing immediate medical or surgical intervention. Patient-side ultrasonography is indicated in all trauma patients to detect internal injury early. If spinal injury is a possibility based on initial assessment, the patient must be immobilized to prevent any deterioration in their neurological condition until further evaluation occurs. Decreased mentation should raise concern for possible traumatic brain injury. Supplemental oxygen and minimal stress are indicated for any patient with respiratory abnormalities in the first instance. Once the patient has been stabilized, a more thorough evaluation of the patient should be performed for all other injuries.

After triage assessment, peripheral intravenous access should be obtained. If this is not possible, a surgical cut-down or intraosseous catheter should be considered. Emergency blood work consisting of a packed cell volume, total solids, blood glucose, lactate and venous blood gas can be taken at the time of catheterization to help guide patient management. Jugular venepuncture or catheterization should be avoided, where possible, in cases of suspected head trauma, to prevent increases in intracranial pressure. Open wounds should be covered in the first instance to reduce the risk of contamination prior to further treatment once the patient is stable. Early opioid analgesia is also paramount for patient welfare and to eliminate any contribution of pain to the cardiovascular assessment. Non-steroidal anti-inflammatories are contraindicated in trauma patients prior to further assessment. Steroid medications should also be avoided due to a lack of proven benefit in this patient population, and increased risk of adverse effects.

The most common cause of cardiovascular instability in the polytrauma patient is hypovolaemic shock secondary to haemorrhage. If hypovolaemia is present, fluid therapy should be started immediately, in the form of either a bolus of 10–20 ml/kg of a balanced isotonic crystalloid (Hartmann's) or 3–5 ml/kg of hypertonic

(7.2%) sodium chloride, given over 15–20 minutes in both cases. Ongoing shock despite aggressive fluid therapy should prompt concern for ongoing bleeding, cardiac arrhythmias, hypoxaemia or organ displacement (herniation). Common sites of internal bleeding include fracture sites and peritoneal, retroperitoneal, pericardial and pleural spaces. In patients refractory to initial crystalloid therapy, it may be necessary to consider an alternative fluid plan, blood transfusion or other means of haemostasis. Frequent patient monitoring is essential owing to the dynamic nature of the trauma patient.

KEY LEARNING OBJECTIVES

- Be able rapidly to identify life-threatening problems commonly seen in polytrauma
- Know how to provide rapid stabilization and appropriate analgesia to the trauma patient
- Be aware of common traumatic injuries in the polytrauma patient and have a strategy for their diagnosis and management

MULTIPLE CHOICE QUESTIONS

1. Which one of the following abnormalities could be consistent with peritoneal bleeding in the trauma patient?
(A) Cullen sign
(B) Normal packed cell volume and a total solids <60 g/l
(C) An increasing volume of peritoneal effusion on serial AFAST (abdominal and focused assessment with sonography for trauma) assessment
(D) Positive glide sign
2. Which one of the following drugs is not an appropriate choice for initial analgesia in a painful trauma patient?
(A) Dexmedetomidine
(B) Methadone
(C) Carprofen
(D) Ketamine
3. Which of the following is the most appropriate fluid resuscitation plan for a dog with moderate hypovolaemic shock and a concurrent traumatic brain injury?
(A) 5 ml/kg hypertonic saline
(B) 1 g/kg mannitol
(C) 20 ml/kg Hartmann's
(D) 15 ml/kg 0.9% sodium chloride

Head trauma

Lindsay Kellett-Gregory

Head trauma in veterinary patients can have many aetiologies. Although ocular, soft tissue and ortho-

paedic injury may occur, TBI may be a significant cause of morbidity and mortality. TBI may be suspected based on neurological findings during initial patient triage, including abnormal mentation, pupil size and reactivity or the presence of other head injury without immediately apparent neurological abnormalities. The Modified Glasgow Coma Score is a useful assessment tool in patients with TBI that are fluid

resuscitated to remove the impact of shock on neurological function.

TBI is divided into primary and secondary injury. Primary injury is that which has already occurred as a direct result of the trauma, such as concussion or contusion. Secondary injury begins minutes after the trauma but may last several weeks and the target for treatment is to minimize neurological injury following trauma. Prevention of hypotension and hypoxaemia are the most important aspects of extracranial stabilization of brain injury.

Intracranial management of brain injury is focused on prevention of an increased intracranial pressure (ICP) that can result in brain herniation and sudden death. Trauma patients are highly susceptible to increases in ICP due to the fixed volume of the skull, with limited space for any oedema or haemorrhage. Severe acute increases in ICP can result in the Cushing reflex, requiring immediate intervention.

There are many aspects to treatment of the patient with severe TBI. Oxygen therapy should be provided to prevent harmful brain hypoxia but should also be titrated to prevent hyperoxia. More aggressive respiratory support may be needed to treat any hypoventilation. Fluid therapy to maintain adequate perfusion is essential, with 0.9% sodium chloride being the isotonic crystalloid fluid of choice at present. Opioid analgesia is also considered an essential part of treatment.

Specific treatment strategies include hyperosmolar therapy to treat intracranial hypertension if present. Mannitol should be given to euvoelaemic patients with increased ICP only and not on a prophylactic basis. Hypertonic saline would be the fluid of choice for patients with evidence of increased ICP and hypovolaemia. Elevation of the head by 15–30 degrees and avoidance of jugular vein compression is also important. There is no evidence that high-dose methylprednisolone is effective. Any seizure activity occurring post TBI requires urgent therapy.

Whilst intracranial imaging has previously been considered for patients with TBI, its use should be reserved for those failing medical management and those for

whom the treatment course will be altered as a result of imaging.

Good supportive and nursing care is essential for a good patient outcome following head trauma. The ultimate prognosis depends on the severity, cause and nature of trauma, as well as the treatment given, but small animals with head trauma have a tremendous ability to compensate and recover from trauma back to a normal quality of life.

KEY LEARNING OBJECTIVES

- Learn how to fully assess the patient with head trauma to detect any traumatic brain injury (TBI)
- Learn about emergency treatment of brain injury
- Understand long-term possible sequelae of head trauma

MULTIPLE CHOICE QUESTIONS

1. Which one of the following is NOT a category assessed in the Modified Glasgow Coma Scale?
 - (A) Brainstem reflexes
 - (B) Motor activity
 - (C) Pain sensation
 - (D) Level of consciousness
2. Which one of the following is NOT a consistent finding seen in the Cushing reflex?
 - (A) Bradycardia
 - (B) Systemic hypertension
 - (C) Decreased mentation
 - (D) Hyperglycaemia
3. Which one of the following is NOT indicated in patients with traumatic brain injury?
 - (A) Methylprednisolone
 - (B) Hyperosmolar therapy
 - (C) Analgesia
 - (D) Intravenous fluid therapy

Uroabdomen: presentation, diagnosis and management

Lindsay Kellett-Gregory

Uroabdomen is defined as leakage of urine into the peritoneal and/or retroperitoneal space, and may occur secondary to trauma, urinary tract obstruction, neoplasia or dehiscence following urinary tract surgery. Consequences include azotaemia, hyperkalaemia, metabolic acidosis, dehydration and, in some cases, shock. These result from retention of potassium and other waste products, in addition to third spacing of fluid into the peritoneal cavity. The chemical peritonitis resulting from

direct contact of urine with the peritoneum may cause acute abdominal pain in some cases.

Clinical signs of uroabdomen are typically non-specific and may include vomiting, anorexia and lethargy. Physical examination findings may include evidence of hypoperfusion, cardiac arrhythmias and abdominal pain. The ability to palpate the bladder does not rule out the possibility of a uroabdomen. Patients with urethral injury may develop inguinal or perineal swelling and extensive bruising.

A diagnosis of uroabdomen is based on a combination of the patient's history and physical examination findings, laboratory assessment of peripheral blood in comparison to abdominal fluid, and imaging findings. A biochemistry panel including electrolytes to detect any hyperkalaemia and azotaemia should be performed in all suspected cases. The degree of hyperkalaemia and azotaemia may be variable, with abnormalities progressing over time. The rise in blood urea nitrogen may be