CLINICAL ABSTRACTS 2021 | ORAL PRESENTATIONS

Following on from the success of BSAVA Virtual Congress 2021, we are delighted to present to you the 2021 Congress Clinical Abstracts. The congress programme committee have spent many hours reviewing some excellent clinical abstracts this year, with the quality and breadth of the 2021 abstract programme a testament to the high level of clinical research that was submitted.

From critical care to endocrinology and from orthopaedics to exotics, the submissions this year have covered a wide range of exciting developments in veterinary medicine, 30 years on from the first BSAVA Congress Abstracts in 1991. As a sign of the changing times since then, a number of the abstracts were presented in online poster form at the 2021 virtual congress, whilst more than 140 will also be presented orally at BSAVA's Congress Clinical Abstract Presentations Online programme on 24th-26th May 2021.

We would like to thank all the submitting authors for their hard work and for choosing BSAVA Congress as a forum in which to share their findings. We are also grateful for the willingness of oral abstract presenters to embrace the challenges of online presentations again this year – hopefully the experience is an enjoyable and rewarding one.

Thank you again for interacting with BSAVA Congress 2021 and see you next year!

Joe Fenn

BSAVA Congress Clinical Abstracts Chair

Note – the abstracts displayed are as supplied and BSAVA has not corrected typographical errors or edited the submission text. Any errors are the responsibility of the authors.

MONDAY 24 MAY 2021

PM Room 2, 12:45 - 18:00 | Primary Practice and the Profession, PetSavers
Student Grants

12:45 Field Efficacy and Safety of a Novel Oral Chewable Tablet Containing Sarolaner, Moxidectin, And Pyrantel Against Naturally Acquired Gastrointestinal Nematode Infections in Dogs Presented as Veterinary Patients in Europe

Csilla Becskei¹, Daphne Fias¹, Sean P. Mahabir², Robert Farkas³

¹Zoetis, Zaventem, Belgium. ²Zoetis, Kalamazoo, USA. ³University of Veterinary Medicine, Budapest, Hungary

Objectives

The efficacy of a novel oral endectocide containing moxidectin, sarolaner and pyrantel was investigated in client-owned dogs with natural infections of *Toxocara canis*, *Toxascaris leonina*, *Ancylostoma caninum* and *Uncinaria stenocephala*.

Methods

Dogs infected with any of the four gastrointestinal nematode species were enrolled in this controlled, masked, randomized study. In total, 194 dogs were treated once orally with the combination of sarolaner (1.2 to 2.4 mg/kg), moxidectin (24 to 48 µg/kg) and pyrantel (5 to 10 mg/kg) and 97 dogs were treated with a positive control product containing afoxolaner (2.50–5.36 mg/kg) and milbemycin oxime (0.50–1.07 mg/kg). Faecal egg counts were conducted before, and 7 days after treatment administration. Efficacy was based on the post-treatment reduction in geometric mean egg counts (per gram faeces) compared to pre-treatment.

Results

Two hundred dogs were infected with *T. canis*, 80 with *A. caninum*, 36 with *U. stenocephala*, and 16 with *T. leonina*. Fifty dogs had mixed infections with two or more species. Post-treatment geometric mean fecal egg counts for *T. canis*, *A. caninum*, *T. leonina*, and *U. stenocephala* relative to pretreatment counts were significantly ($P<0.0001$) reduced by $\geq 98.3\%$ in the group treated with the combination of sarolaner, moxidectin and pyrantel, and by $\geq 97.4\%$ in the afoxolaner and milbemycin oxime-treated group. Both products were well tolerated.

Statement (conclusions)

Oral administration of the combination of 1.2 mg/kg sarolaner, 24 µg/kg moxidectin and 5.0 mg/kg pyrantel provided effective treatment of roundworm and hookworm infection in dogs.

13:00 Raw Diets Safe and Nutritious? U.K. Dog Owner Perspectives on Raw Diets Compared to Cooked Diets

Alysia Empert-Gallegos, Sarah Hill, Philippa S Yam

University of Glasgow School of Veterinary Medicine, Glasgow, United Kingdom

Objectives

What are dog owner perspectives on risks, benefits, and nutritional value of raw diets compared to cooked diets?

Methods

This study is a descriptive, cross-sectional survey targeting Facebook-using pet dog owners in the United Kingdom. Discrete data were analysed using Microsoft Excel. Qualitative data were analysed using RStudio.

Results

The questionnaire was completed by 97 U.K. dog owners; 28.9% fed raw meat-based diets. Participants gave themselves a 4 or 5 out of 5 on dog nutrition knowledge 68.0% of the time, but only gave their veterinarian 4 or 5 on nutrition 46.3% of the time. This dichotomy was more extreme in raw feeders, with only 11.1% giving their veterinarians top scores in nutrition. Raw feeders rated commercial and homemade raw diets as highly nutritious 75.0% and 78.6% of the time, respectively, while only 7.4% rated commercially prepared cooked diets as highly nutritious. Cooked feeders ranked raw diets as highly nutritious less often, but also only ranked cooked diets as highly nutritious 60.9% of the time. All participants agreed that cooked diets were low risk to human health. Raw feeders ranked raw diets as highly risky to human or dog health 3.6% of the time in all categories but deemed cooked diets risky to dog health 53.6% of the time. This work was supported by MSD Animal Health.

Statement (conclusions)

This research bridges the gap of knowledge about owner perspectives around feeding raw and will help veterinarians educate and converse with clients about making the healthiest and safest diet choice for their dogs.

13:15 Evaluation of A Point-Of-Care Dot Enzyme-Linked Immunosorbent Assay to Detect Antibodies Against Canine Parvovirus, Distemper Virus and Adenovirus Before Revaccination.

Ditte Erika Leth Vasby, Line Kristensen, Agnethe Wallin, Jo Fjeldsted-Holm Lundsgaard, Tina Møller Sørensen

Department of Veterinary Clinical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Frederiksberg C, Denmark

Objectives

Antibody measurement is the new recommended practice to evaluate the individual need for canine core vaccination boosters, according to the World Small Animal Veterinary Association (WSAVA).

The objectives are to evaluate the accuracy of a point-of-care titer test to detect canine antibodies in primary practice, and if accuracy improves when applying WSAVA recommendations regarding antibody cut-off values for gold standard analysis.

Methods

A prospective non-inferiority study (margin of 15%) of a point-of-care semi-quantitative titer test was performed on canine serum samples (n=82) and compared to gold standard analysis. Virus neutralization (VN) was performed for canine distemper virus (CDV) and canine adenovirus (CAV) while haemagglutination inhibition (HI) was performed for canine parvovirus (CPV) as quantitative gold standard. Sensitivity, specificity, positive and negative predictive values of the point-of-care titer test were determined for gold standard titer cut-off ≥ 16 , ≥ 32 and ≥ 80 for CAV, CDP and CPV, respectively. For WSAVA cut-offs all titers ≥ 10 was considered positive.

Results

The titer test showed a sensitivity of 91% [84;96], 87% [78;93] and 100% [96;100], and a specificity of 100% [37;100], 52% [33;71] and 100% [5;100] for CAV, CDV and CPV, respectively. No significant difference was seen when applying WSAVA cut-offs. As specificity is considered the most important parameter, the titer test was considered inferior to gold standard VN with respect to CDV antibodies.

Statement (conclusions)

Care should be taken when implementing and interpreting point-of-care titer tests in primary practice. More emphasis on high specificity is needed when developing and refining new in-house tests for antibody titer evaluation.

13:30 Cooling Methods Used in Dogs with Heat-Related Illness Under UK Primary Veterinary Care During 2016-2018.

Emily Hall¹, Anne Carter¹, Jude Bradbury¹, Dominic Barfield², Dan O'Neill²

¹Nottingham Trent University, Nottingham, United Kingdom. ²Royal Veterinary College, London, United Kingdom

Objectives

Greater duration and extent of hyperthermia in heat-related illness cases is associated with poorer prognosis. Rapid, effective cooling is critical for improved outcomes. Water immersion and water spray with air movement are currently considered the most effective cooling methods. This study explored cooling methods used in dogs with heat-related illness presenting for first-opinion veterinary care, and compared utilised cooling practices to best practice.

Methods

Dogs with heat-related illness events were identified from VetCompass clinical records. Clinical severity and information on cooling methods pre/post presentation were extracted. Associations between event severity and cooling methods used were assessed using a chi-squared test.

Results

The analysis included 810 heat-related illness events. No active cooling was recorded in 47.9% (388/810 events). Among 422 (52.1%) actively cooled dogs, cooling methods were: wet towels (35.3%), water spray/water immersion (33.2%), air movement (22.5%), ice/ice packs (11.8%), cold

intravenous-fluids (5.9%), alcohol on footpads (5.2%), cold-water enema (1.9%) and unspecified (28.7%). Multiple cooling methods were used in 34.1% of events, with wet towels plus air movement (8.5%) the most common combination reported.

Compared to mild and moderate cases, severe cases were three times more likely to receive cold intravenous-fluids ($p=0.003$), and six times more likely to receive cold-water enemas ($p=0.007$).

Statement (conclusions)

These findings benchmark active cooling actions used to manage heat-related illness in dogs in the UK; the use of active cooling was not documented in almost half of cases. These results suggest the value of improved public and veterinary education to increase use of effective, evidence-based cooling.

13:45 VetCompass Clinical Grading Tool for Heat-Related Illness in Dogs – A Novel Tool to Support Clinical Decision-Making in Primary-Care Practice.

Emily Hall¹, Anne Carter¹, Jude Bradbury¹, Dominic Barfield², Dan O'Neill²

¹Nottingham Trent University, Nottingham, United Kingdom. ²Royal Veterinary College, London, United Kingdom

Objectives

Historically, classification of heat-related illness in dogs relied on body temperature and neurological evaluation, extrapolated from traditional human classifications. Novel systems in human medicine now embrace the multi-systemic progressive nature of heat-related illness. This study aimed to adapt these novel human systems for use in dogs and explore the resultant predictive value for primary-care triage.

Methods

Dogs with heat-related illness were identified from VetCompass UK primary-care clinical records. Clinical presentation and outcome data were extracted, events were then retrospectively graded as mild, moderate or severe using a novel clinical grading tool which utilised only clinical signs. The ability of the grading tool to predict fatality was explored using logistic regression.

Results

The study included 856 heat-related events (2016-2018). The most frequent clinical signs were altered respiration (63.7%) and lethargy (44.3%). Clinical signs associated with increased risk of death were: abnormal mentation including coma (x13.3), stupor (x9.6), multiple seizures (x6.4), gastrointestinal haemorrhage (x5.2), petechiae/purpura (x4.2) and ataxia (x3.3).

The survival rate for cases graded by the tool as mild (altered respiration, lethargy, episodic collapse), moderate (gastrointestinal signs, a single seizure) and severe (abnormal mentation, gastrointestinal haemorrhage, petechiae/purpura) were 97.8%, 94.5% and 43.2% respectively.

Severe events had 58.1 (95%CI 25.1-134.4) times the odds of death compared to mild events.

Statement (conclusions)

The marked survival difference between grades highlights the prognostic value of the novel clinical grading tool in practice. The VetCompass grading tool used in combination with patient signalment and history could improve clinical decision making for dogs with heat-related illness.

14:30 The Scoop on Puppies That Eat Their Own Poop: Is Autocoprohagia Retained as Puppies Age?

Rachel Kinsman¹, Rachel Casey¹, Toby Knowles², Séverine Tasker^{3,2}, Joshua Woodward¹, Rosa Da Costa¹, Sara Owczarczak-Garstecka¹, Jane Murray¹

¹Dogs Trust, London, United Kingdom. ²Bristol Veterinary School, Bristol, United Kingdom. ³Linnaeus Group, Shirley, United Kingdom

Objectives

To investigate if autocoprohagia is retained as puppies age, and, for puppies displaying the behaviour at age 16-weeks, if owner response is associated with the occurrence of autocoprohagia at 7-months.

Methods

In an ongoing UK/ROI canine longitudinal study, owner-completed survey data about autocoprohagia (when puppies were aged 16-weeks and 7-months) were analysed. The prevalence of autocoprohagia was calculated for both timepoints, and the McNemar test was used to test for a change in prevalence between the timepoints. A Chi-squared test was used to test for association between owner-reported responses to seeing autocoprohagia at age 16-weeks and presence/absence of autocoprohagia at age 7-months.

Results

At age 16-weeks and 7-months, 11.7% (257/2200) and 4.6% (69/1515) of puppies, respectively, had reportedly eaten their own faeces in the last seven days. Autocoprohagia prevalence decreased between the two timepoints (n=1,349), p<0.001.

Autocoprohagia at 7-months was reported for 22.5% (9/40) of puppies whose owners had reported using only non-aversive responses (clearing up the faeces, calling dog away, distracting using a toy/treat, ignoring dog) at 16-weeks, compared with 30.6% (34/111) whose owners reported using either only aversive responses (use of loud/sudden noise, telling off, pulling away, putting on lead, rubbing dog's nose in it), or both aversive and non-aversive responses. The association between owner response to autocoprohagia at 16-weeks (n=151) and subsequent autocoprohagia did not reach statistical significance, p=0.329.

Statement (conclusions)

Autocoprohagia prevalence decreased as puppies aged. This analysis will be repeated with a larger sample size and greater age span to better understand this behaviour.

14:45 What Makes Dogs Tick? Risk Factors and Clinical Management of Tick Infestations in UK Dogs

Rinrada Komutrattananon¹, Ashley Hartley², Camilla Pegram¹, David Church¹, Dave Brodbelt¹, Dan G. O'Neill¹

¹Royal Veterinary College, Hertfordshire, United Kingdom. ²College of Veterinary Medicine, University of Tennessee, Knoxville, USA

Objectives

To report prevalence, risk factors, clinical management, and outcomes of tick infestation in UK dogs.

Methods

VetCompass cohort study using 905,554 dogs under veterinary care in 2016. Tick infestation cases were identified using electronic searches and compared to dogs under veterinary care for other reasons during 2016. Risk factor analysis used multivariable logistic regression modelling.

Results

One-year (2016) period prevalence of tick infestation was 0.5% (95%CI 0.2-0.7). Medium-haired dogs were predisposed (OR 2.3, 95%CI 2.0-2.6) compared with short-haired dogs. Cairn Terriers (OR 3.2, 95%CI 1.4-7.0) had the highest odds while Staffordshire Bull Terriers (OR 0.3, 95%CI 0.2-0.5) had the lowest compared to Labrador Retrievers. Cases occurred in every season, but predominantly summer (52.7%) and spring (27.0%). Ticks mainly in the head area (41.8%), especially around the eyes (27.9%) and ears (26.7%), followed by forelimbs (8.1%) and neck (6.2%). Of the tick cases, 12.4% of the wounds were infected. Most ticks were removed by clinical staff (58.7%). Medical therapy was prescribed in 51.5% of cases. Of those prescribed medical therapy, 56.1% cases were prescribed acaricides. NSAIDs, glucocorticoid or/and antibiotic were prescribed to 6.3% of tick cases.

Statement (conclusions)

This national tick infestation prevalence benchmark of 0.5% in 2016 can help with tick surveillance to assess trends over time, especially in response to changing acaricide preventative use and global warming. Veterinarians could prioritise routine tick surveillance especially in medium-haired dogs, and around the eyes and ears. Acaricides and vigilant grooming for ticks should be prioritised during summer months.

15:00 Concerns and Experiences of Accessing Veterinary Care During the COVID-19 Pandemic: A Mixed-Methods Analysis of Dog Owners' Responses.

Sara C. Owczarczak-Garstecka, Katrina E Holland, Katharine L. Anderson, Rachel A. Casey, Robert M. Christley, Lauren Harris, Kirsten M. McMillan, Rebecca Mead, Jane K. Murray, Lauren Samet, Melissa M. Upjohn

Dogs Trust, London, United Kingdom

Objectives

To explore dog owners' concerns and experiences related to accessing veterinary care during the UK's COVID-19 pandemic.

Methods

Data were obtained through two owner-completed surveys conducted in May 2020 (during the first nationwide lockdown) and during October 2020 and an electronic diary completed by owners (April-May 2020). Responses to three open-ended questions and electronic diaries were analysed qualitatively to identify key themes. Responses for questions concerning access to veterinary care from the second survey were summarised with descriptive statistics.

Results

During the lockdown, availability of veterinary care worried 32.4% (n=1,431) of respondents. However, 98.1% (n=1,745) of those who needed to access a veterinarian reported being able to do so. Significant qualitative themes were: Concerns around access to a veterinarian, delays in treatments and worries about the potential impacts of these delays on dogs' health. In relation to experiences since 23/03/2020, in the October survey, 30.4% (n=37/127) of dogs that owners planned to neuter were not yet neutered and neutering of further 16.5% (n=21/127) was delayed. Of dogs that owners intended to vaccinate, 11.0% (n=110/1,000) were not yet vaccinated and an additional 23.4% (n=234/1,000) experienced a delay in vaccination.

Respondents were also concerned about being unable to accompany dogs into the veterinary practice, especially if they cared for dogs with behavioural issues or elderly dogs, with fears around the potential for euthanasia in the owner's absence.

Statement (conclusions)

Although data collection is ongoing, preliminary results highlight that COVID-19 restrictions have delayed prophylactic treatments for some dogs and caused worry for about a third of owners. Our findings can inform veterinarians' dog owner engagement to address their concerns.

15:15 Euthanasia in Dogs in the UK; What Drives Decision-Making?

Camilla Pegram¹, Carol Gray², Rowena Packer¹, Ysabelle Richards¹, Dave Brodbelt¹, David Church¹, Dan O'Neill¹

¹Royal Veterinary College, Hatfield, United Kingdom. ²University of Liverpool, Liverpool, United Kingdom

Objectives

Veterinarians describe euthanasia as "the best and the worst" of their clinical role. This study explores how dogs die in UK general practice, and what demographic and clinical factors are associated with euthanasia relative to unassisted death.

Methods

Deceased dogs during 2016 were identified from anonymised clinical records in VetCompass. Deaths were categorized as euthanasia or unassisted. Demographic and cause of death data were extracted. Risk factor analysis for euthanasia relative to unassisted death used multivariable logistic regression.

Results

From 29,163 deceased dogs, 26,676 (91.5%) were euthanased and 2,487 (8.5%) died unassisted. Using neoplasia as the baseline, 6/20 disorder groups had higher odds in euthanased dogs. The disorders that were most likely to end in euthanasia compared with unassisted death were poor quality of life (odds ratio [OR] 15.64), undesirable behaviour (OR 13.10) and spinal cord disorder (OR 6.24). Conversely, traumatic injury (OR 0.06), complication associated with clinical care (OR 0.09) and heart disease (OR 0.13) were least likely to end in euthanasia. Bulldogs (0.38), Pugs (0.45), and West Highland White Terriers (0.60), had reduced odds of euthanasia compared with Labrador Retrievers. No breeds had significantly increased odds of euthanasia. Dogs aged ≥ 15 years had 8.33 times the odds of euthanasia compared with dogs < 6 years. Bodyweight ≥ 30 kg had 1.32 times the odds of euthanasia compared with dogs < 10 kg.

Statement (conclusions)

These findings can support veterinarians undertaking euthanasia conversations with clients, by highlighting factors that may influence such decisions and offering a basis for shared decision-making based on evidence.

15:45 Owner Strategies for Vaccination and Revaccination in Pedigree and Non-Pedigree Dogs

Megan Robson, Mark Dunning

University of Nottingham, Nottingham, United Kingdom

Objectives

Identifying motives behind owners' vaccination strategies and why they choose not to vaccinate or revaccinate their dogs to aid the discussions on how to improve the overall vaccination uptake in the UK.

Methods

A survey was constructed and distributed via breed groups and social media. The survey ran for 23 days, with 286 responses (totalling 790 dogs, of which 677 were pedigree and 113 non-pedigree). The data was analysed and qualitative answers filtered to find common themes. A Chi-squared test was used to determine any significant differences between datasets (significance set at $P < 0.05$).

Results

A total of 4.6% of pedigree and 8% of non-pedigree dogs received no vaccinations, a further 33.1% pedigree and 29.2% non-pedigree dogs were not revaccinated after their primary vaccination course. Leptospirosis uptake was particularly poor with only 68.7% pedigree and 67.3% non-pedigree dogs receiving vaccinations. Popular reasons against vaccinating included a preference to titre test, not believing they are necessary, cost and preference for homeopathy. Breeds standing out with lower uptake: 30.5% of Huskies were unvaccinated and 49.4% of Irish Wolfhounds were not revaccinated.

Statement (conclusions)

The results indicate that UK dogs may not be sufficiently vaccinated to provide adequate herd immunity and many may be at risk of disease, especially leptospirosis. Further work reviewing the validity of titre testing and more detailed information in selected breeds regarding vaccinations will aid efforts in improving vaccination uptake in UK dogs.

16:00 Characterising First-Line Empirical Treatment for Coughing - A Retrospective Evaluation Of 83 Dogs.

Joshua Hardwick^{1,2}, Jennifer Reeve¹

¹University of Bristol, Bristol, United Kingdom. ²Langford Vets, Bristol, United Kingdom

Objectives

Primarily, to characterise prescribing tendencies of primary care veterinary surgeons for dogs presenting with a cough. Secondly, to report the definitive diagnosis in this cohort and correlate the class(es) of drug(s) prescribed at initial presentation with the final diagnosis – specifically the correlation between antimicrobial use and diagnosis of an infectious cause of the cough.

Methods

Retrospective review of case records of dogs presenting to a referral centre between 2015-2020, following initial presentation to their primary veterinary clinic with coughing as a primary complaint. Cases were excluded if clinical histories were not available for review or a final diagnosis was not reached at referral. Included cases were stratified into the following categories based upon the final diagnosis (a) confirmed infectious (b) suspected infectious (c) non-infectious. Descriptive statistics were used to compare the use of different drug classes across these groups.

Results

83 dogs met the inclusion criteria of which 15 (18.1%) had a confirmed infectious cause, 6 (7.2%) had a suspected infectious cause and 62 (74.7%) had a non-infectious cause. Initial treatment comprised antimicrobials in 43 dogs (51.8%), non-steroidal anti-inflammatory drugs in 41 (49.4%), corticosteroids in 3 (3.6%), anthelmintics in 5 (6%) or drugs of a different class in 9 (10.8%).

Statement (conclusions)

In this study, 21 dogs (25.3%) had a confirmed or suspected infectious cause for their cough, despite over half (n=43, 51.8%) having received antimicrobials empirically at first presentation. Further studies identifying presenting features that would facilitate identification of dogs who would benefit from antimicrobials on first presentation would be valuable.

16:15 Vulval Conformation in The Entire Bitch and The Effects of Pre And Post-Pubertal Neutering on Vulva Development and Urogenital Disease.

Jasmine Godfrey-Hunt¹, Caitlin Pagett¹, Rachel Moxon², Gary England¹

¹University of Nottingham, Nottingham, United Kingdom. ²Guide Dogs Breeding Centre, Warwickshire, United Kingdom

Objectives

- 1) Describe normal vulval appearance for entire adult bitches, and compare to bitches neutered pre-pubertally and post-pubertally
- 2) Examine associations between vulval development and appearance and urogenital disease

Methods

Digital images of the vulva and vulval measurements were analysed from 412 adult large-breed bitches (106 entire, 155 neutered pre-pubertally and 151 neutered post-pubertally). Vulva size (cm) and development, vulval dorsal skin fold characteristics, recessed and juvenile vulval appearance were described and compared between the groups. Retrospective examination of health records identified cases of urogenital disease.

Results

Experiencing puberty resulted in greater vulval development; pre-pubertally neutered bitches had significantly smaller vulvas than post-pubertally neutered and entire bitches ($P < 0.001$). There was no difference in vulva length and width between bitches neutered post-pubertally and entire bitches. Recessed and juvenile vulvas were more prevalent in pre-pubertally neutered bitches at 17-months of age (12.9% and 15.0% respectively, compared to 2.9% and 1.4% for post-pubertally neutered bitches). Prevalence of UTI was greater in bitches neutered pre-pubertally (9.0% compared to 2.0%; $P = 0.007$).

Statement (conclusions)

Neutering and experiencing puberty impacts vulval development and subsequent predisposition to urogenital disease in bitches, although the incidence of disease was low. Examination of the vulva could be considered as part of routine decision-making regarding neutering.

16:30 Journal Advertising: What Does It Tell Us?

Natasha Basham¹, Simran Floyd¹, Lisa Morrow^{2,1}, Rachel Dean^{3,1}, Marnie Brennan^{2,1}

¹The University of Nottingham, Nottingham, United Kingdom. ²CEVM, Nottingham, United Kingdom.

³Vet Partners, York, United Kingdom

Objectives

Previous medical research indicates that, while advertisements in medical journals can have a profound impact on clinician prescribing habits, they often contain substandard provision of peer-

reviewed references. Minimal research exists exploring this concept within the veterinary industry. Therefore, the aim of this study was to interrogate advertisements in 2 veterinary periodicals.

Methods

All advertisements published in the Veterinary Times and Veterinary Record over a 1-year period that fit the inclusion criteria were identified. Characteristics such as target species, product action, company, size of advert, and type of references supplied were assessed. Two reviewers independently extracted and compared the data.

Results

A total of 524 adverts which met the inclusion criteria were identified; 16.1% of these contained no references at all. Overall, 882 references mentioned in the adverts were analysed, of which only 111 (12.6%) were peer-reviewed. Preliminary findings suggest associations between factors such as target species and product action on whether peer-reviewed references were provided.

It appears that peer-reviewed references are not often provided in adverts. This has implications for veterinary professionals being able to apply the principles of evidence based veterinary medicine into clinical practice. Future work should investigate the practical consequences of these findings.

Statement (conclusions)

This work highlights that references in adverts are frequently not peer-reviewed which is a potential barrier to practitioners when making evidence-based decisions in clinical practice. The findings provide a starting point for discussion within the veterinary industry concerning advert presentation and draws attention to current issues veterinary professionals should be aware of.

16:45 Surgical Safety Checklists in Veterinary Practice: Current Implementation Within UK Practices and Attitudes Towards Their Use

Jessica Hill^{1,2}, Gemma Irwin-Porter¹, Louise Buckley³

¹University of Bristol Veterinary School, Bristol, United Kingdom. ²Paragon Veterinary Referrals, Wakefield, United Kingdom. ³University of Edinburgh, Edinburgh, United Kingdom

Objectives

This study had two key objectives:

- To understand how respondents' veterinary practices (VPs) used surgical safety checklists (SSCs)
- To identify risk factors for poorer respondent attitude towards SSCs (all respondents) or VPs not using SSCs (qualified respondents)

Methods

A prospective observational cross-sectional survey with convenience sampling was utilised. From this, a summative composite attitudinal scale (13 statements; Cronbach alpha: 0.932; scale: 0 – 39,

39 = most positive) was developed. Risk factors were analysed using Generalised Linear Models, reported using means or odds ratios, with 95% confidence intervals.

Results

Of 517 respondents, 70.0% worked in practices using SSCs. Of these, 87.0% always used SSCs, 19.2% adapted SSCs to suit procedures and 60.2% had an SSC Standard Operating Procedure.

Attitudes to SSCs were generally favourable (mean: 28.3, $\pm$ S.D. 6.8). A more positive respondent attitude was associated with: working in a VP employing at least one RVN with a post-qualifying qualification ($p = 0.027$), using SSCs ($p = 0.001$), working in a training practice ($p = 0.016$), undertaking self-directed reading ($p = 0.025$), or completing relevant CPD post-qualification ($p = 0.033$) around SSCs.

Risk factors for VPs not using SSCs included: Practice Standards Scheme (PSS) non-membership (OR 2.0, 1.1. – 3.4), not having RCVS hospital status (OR 1.9, 1.1. – 3.5) or being a mixed first opinion VP (OR 2.4, 1.2 – 5.0).

Statement (conclusions)

Encouragingly, most VPs utilised SSCs. The findings suggest familiarity, education and the RCVS PSS improve uptake and attitudes. Further research should identify mixed practice-specific barriers to SSC use.

17:00 Prevalence of Alimentary and Respiratory Disease in Brachycephalic Dog Breeds Presenting to Primary Care Practice.

Will Petchell¹, Rachel Burrow², Peter-John Noble¹, Harriet Broome¹, Marisol Collins¹

¹University of Liverpool, Liverpool, United Kingdom. ²Northwest Veterinary Specialists, Runcorn, United Kingdom

Objectives

The aims were to measure prevalence of alimentary and respiratory disease in French Bulldogs (FBD), English Bulldogs (BD) and Pugs attending first-opinion practice and to identify differences between breeds to improve management of brachycephalic dogs.

Methods

Clinical records of 260 FBD, 238 BD and 252 Pugs attending over 500 first-opinion practices from October 2014 - May 2020 from the Small Animal Veterinary Surveillance Network (SAVSNET) database were analysed. Inclusion criteria included patient age of <1 year at initial presentation and >36 months between the first and last clinical entry. Clinical signs of oesophageal, gastro-oesophageal, gastric, lower gastrointestinal, upper respiratory tract, lower respiratory tract and miscellaneous (exercise intolerance and syncope) were recorded as present or absent. Data were stored and analysed in Microsoft Excel 2015 (Microsoft Corporation).

Results

FBD had the highest prevalence of alimentary tract disease while Pugs had the highest prevalence of respiratory tract and miscellaneous disease. There were significant differences in the prevalence of oesophageal, gastric and lower gastrointestinal clinical signs between breeds ($p < 0.017$). FBD had significantly increased odds ratios of having oesophageal (2.61), gastric (1.86) and lower gastrointestinal disease (1.61). Pugs had significantly increased odds ratios of having upper respiratory tract disease (1.61).

Statement (conclusions)

Our results show important differences in the prevalence and likelihood of alimentary tract and respiratory disease between brachycephalic breeds attending primary care vets. The difference in clinical sign localisation and known association between gastro-oesophageal and obstructive airway diseases highlights the importance of considering the contribution of gastroenteric disease to clinical presentation when treating FBD in general practice.

17:15 Retrospective Study of Feline Lymphoma: Distribution, Clinical Signs, Nervous System Involvement and Diagnostic Testing (2013-2019) - Alexander Davies

Alexander M. Davies, Elsa Beltran, Irina Gramer

Royal Veterinary College

Objectives:

The aim of this study was to assess the relative frequency of different forms of feline lymphoma, especially those involving the nervous system (NS), and review their clinical features and diagnostic findings.

Methods:

Medical records of cats referred to the Queen Mother Hospital for Animals between 2013 and 2019 were reviewed. Cats were included if a lymphoma diagnosis was confirmed with cytology and/or histopathology. Signalment, retroviral status, tumour anatomical location, clinical signs at presentation, diagnostic testing and staging results were collected.

Results:

Two hundred and fifty-five cats were included. Median age at presentation was 9.7 years. One hundred and sixty-three were males and ninety were females. Seventy-six cats received retroviral testing, 5.3% were positive for FeLV and 13.2% positive for FIV. Anatomical locations of lymphoma included alimentary (55.7%), hepatic (14.5%), splenic (9.4%), mediastinal (10.6%), nasal (8.2%), cutaneous (2.4%), multicentric (19.6%), renal (14.5%) and NS (9.4%). Cats with NS lymphoma presented with chronic (79.2%), progressive (100%), lateralised (58.3%) and non-painful (58.3%) neurological deficits. The lesions affecting the NS were seen with advanced imaging in seventeen cats. Alimentary lymphoma (32.2%) and renal lymphoma (25.8%) were most frequently seen in cats with neurological deficits.

Impact/Clinical Significance:

NS lymphoma is relative infrequent compared to alimentary, mediastinal, multicentric and cutaneous forms, however it must be considered when compatible clinical signs are present. The clinical and diagnostic data shown in this study contributes to the clinical decision-making in primary care and referral practice.

Funding/Declarations of interest:

This project was funded with a PetSavers Student Research Project Grant.

Ethics declaration:

The authors declare no conflicts of interest. The work described in the abstract complies with UK legislation pertaining to animal welfare, ethics and data protection. This abstract will not be presented or published elsewhere before BSAVA Congress 2021. This abstract submission has been approved by all authors.

17:45 Radiographic Assessment and Observer Reliability in Cases of Implant Loosening Following Canine Total Hip Replacement (THR)

Samuel Pearce¹, Eithne Comerford², Sumaya Allaith², Sorrel Langley-Hobbs³

¹Bristol Veterinary School, ²University of Liverpool, ³Bristol Veterinary School,

Objectives:

To describe common radiographic features in cases of septic and aseptic loosening of canine THR implants and measure inter- and intra-observer reliability for radiographic assessment of loose THR implants.

Methods:

A retrospective study was performed using cases of implant loosening identified from the University of Liverpool – British Veterinary Orthopaedic Association Canine Hip Registry (UoL-BVOA CHR) and University of Bristol - Langford Vets. All cases of septic or aseptic loosening were included. Four board-certified orthopaedic surgeons reviewed anonymised affected radiographs on two occasions, according to criteria about bone changes, implant position, radiolucency changes, surgical technique and whether they thought the case was septic or aseptic. Reliability was assessed using Krippendorff's alpha.

Results:

Eighty-eight cases met the inclusion criteria. Clinical records and radiographs were obtained for thirty-three cases. In the first review, notable reported features included periosteal reaction (75.8% of cases), femoral remodelling (72.0%), acetabular remodelling (48.5%), implant subsidence (25.0%) and implant rotation (18.9%). Surgeons deemed the stem sizing and positioning post-THR were correct in 71.2% and 59.1% of cases, respectively. However, these results were inconsistent with the second review. The orthopaedic surgeons correctly identified 27.3 - 66.7% of cases as septic or aseptic. Inter-rater reliability was 0.03 – 0.5 for each variable in the first round and all were <0.16 in the second round.

Impact/Clinical Significance:

Septic and aseptic loosening can be difficult to diagnose and differentiate between from radiographs alone. These results suggest that even experienced observers have low agreement when assessing radiographs of THR implant loosening, but further research is needed to confirm this.

Funding/Declarations of interest:

This project was funded with a PetSavers Student Research Project Grant.

Ethics declaration:

The authors declare no conflicts of interest. The work described in the abstract complies with UK legislation pertaining to animal welfare, ethics and data protection. This abstract will not be presented or published elsewhere before BSAVA Congress 2021. This abstract submission has been approved by all authors.